

Variations in Carbon Dioxide Gas Exchange and Productivity of the Major Crops of the Haeen Catchment, South Korea



University of Bayreuth



Steve Lindner, Bora Lee, Peng Zhao, Janine Kettering, Hyojung Kwon, Dennis Otieno and John Tenhunen

Contact: Steve.Lindner@yahoo.de

Introduction:

The Asian region is a major contributor to global atmosphere / biosphere exchange of energy, water, carbon dioxide and other trace gases. Furthermore, it accounts for ca. 20% of the world's agricultural land (Fan 2005). Agricultural production over large parts of Asia and under the common farming practices carried out in South Korea is input-intensive with respect to fertilizer, pesticides, and machinery use. The objectives of the current study were to evaluate the outputs from such management practices, e.g., to:

1. Quantify the diurnal and seasonal CO₂ exchange of the dominant agricultural crops within the Haeen landscape,
2. Quantify biomass production and its partitioning in the major crops throughout the growing season,
3. Identify key drivers of agro-ecosystem CO₂ exchange and productivity, and
4. Relate plot level CO₂ exchange to landscape level exchange fluxes, biomass development and total farm yields.

Hypothesis: Yield of the agricultural crops raised in Haeen Catchment is determined by seasonal changes in carbon uptake capacity, the duration of carbon uptake and the patterns occurring in carbon partitioning.

Materials and Methods:




Figure 1: A) Experimental set up for the CO₂ chamber measurements in the crop fields. B) Installed light chamber during the CO₂ exchange measurements in the rice field. C) Top down detail picture of a rice plot.

Net ecosystem CO₂ exchange (NEE) and ecosystem respiration (R_{eco}) of the dominant crops in the Haeen Catchment were observed during 2009 and 2010 with a systematic rotation over 6 plots (Figure 1A) using manually-operated, closed light (see Figure 1B and 1C) and dark gas exchange chambers (cf. Otieno et al. 2012). A set of 6 soil frames were inserted 20 cm into the ground as bases for CO₂ measurement chambers. After CO₂ measurements, all of the biomass within each of the 40 cm by 40 cm soil frames was harvested to determine dry weight of plant organs and leaf area. The samples were also analyzed for N and C contents. Average yield of the catchment was estimated from harvests in 32 randomly selected fields distributed over the entire catchment. Empirical description of the measured NEE fluxes was performed with a non-linear least squares fit of the data to a hyperbolic light response model (Owen et al. 2007).

Results and Conclusions:

Table 1: Dry weight of biomass in 2009

	Day of Year 2009	Leaves		Stems		Root		Grains/Tuber Dry	
		Dry weight [kg/m ²]	SD	Dry weight [kg/m ²]	SD	Dry weight [kg/m ²]	SD	Dry weight [kg/m ²]	SD
Radish	163	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	176	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	196	0.04	0.00	0.02	0.00	0.03	0.00	0.00	0.00
	213	0.07	0.01	0.06	0.01	0.24	0.03	0.00	0.00
	247	0.11	0.02	0.07	0.01	0.32	0.08	0.00	0.00
	253	0.06	0.01	0.07	0.02	0.35	0.07	0.00	0.00
Cabbage	148	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	162	0.04	0.02	0.02	0.01	0.00	0.00	0.00	0.00
	177	0.16	0.05	0.11	0.04	0.01	0.01	0.00	0.00
	197	0.27	0.05	0.26	0.06	0.02	0.01	0.00	0.00
	159	0.10	0.03	0.05	0.02	0.03	0.02	0.09	0.04
	174	0.08	0.06	0.07	0.07	0.02	0.01	0.32	0.21
Potato	189	0.09	0.03	0.04	0.02	0.03	0.01	0.62	0.18
	220							1.44	0.22
	147	0.01	0.00	0.00	0.02	0.00	0.00	0.00	0.00
	155	0.00	0.00	0.00	0.03	0.03	0.03	0.00	0.00
	156	0.01	0.00	0.00	0.00	0.01	0.01	0.00	0.00
	164	0.02	0.00	0.01	0.01	0.03	0.00	0.00	0.00
Rice	175	0.05	0.01	0.01	0.01	0.16	0.11	0.00	0.00
	203	0.27	0.05	0.37	0.12	1.17	0.54	0.00	0.00
	260	0.25	0.02	0.84	0.15	0.54	0.08	0.91	0.10
	169	0.01	0.01	0.01	0.00	0.01	0.00	0.00	0.00
	179	0.03	0.01	0.02	0.01	0.01	0.00	0.00	0.00
	202	0.11	0.06	0.12	0.07	0.03	0.01	0.00	0.00
Bean	235	0.35	0.06	0.49	0.14	0.07	0.02	0.10	0.03




Figure 2: Seasonal course of Reco and NEE in (A) 2009 and (B) 2010.




Figure 3: Fitted light curves from the daily courses for CO₂ exchange measurements with (C) potato, (D) rice and (E) radish in 2010.

Results of CO₂ exchange as observed during 2009 and 2010 are shown in Figure 2 A and B and biomass development in Table 1 for the most important annual dryland crops (potato, radish, cabbage, bean) and rice in the Haeen Catchment. The main determinants of CO₂ exchange and biomass development in the catchment were radiation and temperature. Peak rates of NEE and GPP during the season were dependent on the developmental stages of the crops, daily flux rates were influenced by the time course for PAR (see Fig. 3). Local farmers played a strong role by determining the planting dates. Upscaling from plot to landscape level is planned, using eddy covariance data and the PIXGRO-model (Lee et al. 2010).

References:

Fan S, Kang CC 2005: Is small beautiful? Farm size, productivity, and poverty in Asian agriculture. *Agricultural Economics* 32, 135-146.
 Lee B, Tenhunen JD, Geyer R, Seo B, Li Y, Kang S 2010 Landscape level carbon and water balances and agricultural production in mountainous terrain of the Haeen Basin, South Korea., *Proceedings of the Symposium: LANDMOD2010 International Conference on Integrate Landscape Modelling, Montpellier Feb. 3-5, 2010*. Internet publication: www.symposcience.org
 Otieno D, Lindner S, Muhr J, Borken W 2012: Sensitivity of peatland herbaceous vegetation to vapor pressure deficit influences on net ecosystem CO₂ exchange. *Wetlands* 32, 895-905.
 Owen, K.E. et al. 2007: Comparison of seasonal changes in CO₂ exchange capacity of ecosystems distributed along a north-south European transect under non-water-stressed conditions. *Global Change Biol.* 13, 734-760.